



Binge Alcohol Exposure Triggers Atrial Fibrillation Through T-Type Ca^{2+} Channel Upregulation via Protein Kinase C (PKC) / Glycogen Synthesis Kinase 3β (GSK 3β) / Nuclear Factor of Activated T-Cells (NFAT) Signaling — An Experimental Account of Holiday Heart Syndrome —

Yan Wang, MD, PhD; Masaki Morishima, PhD; Dan Li; Naohiko Takahashi, MD, PhD;
Tetsunori Saikawa, MD, PhD; Stanley Nattel, MD, PhD; Katsushige Ono, MD, PhD

Background: The association between binge alcohol ingestion and atrial fibrillation (AF), often termed “holiday heart syndrome”, has long been recognized. However, the underlying cellular and molecular mechanisms are unknown.

Methods and Results: An experimental model of binge alcohol-induced AF was developed to elucidate the mechanisms linking acute ethanol exposure to changes in ion channel transcription and AF susceptibility. AF-susceptibility during transesophageal electrical stimulation was enhanced 8 h after, but not immediately or 24 h after, acute alcohol intake. T-type calcium channel (TCC) blockade and calcineurin inhibition diminished the AF-promoting effect of ethanol. Long-term (8–24 h) exposure to ethanol augmented TCC isoform-expression (Cav3.1 and Cav3.2) and currents in cardiomyocytes, accompanied by upregulation of the transcription factors, Csx/Nkx2.5 and nuclear factor of activated T-cells (NFAT), in the nucleus, and of phospho-glycogen synthesis kinase 3β (GSK 3β) in the cytosol. Inhibition of protein kinase C (PKC) during the 7- to 8-h period following ethanol exposure attenuated susceptibility to AF, whereas acute exposure did not. GSK 3β inhibition itself upregulated TCC expression and increased AF susceptibility.

Conclusions: The present study results suggest a crucial role for TCC upregulation in the AF substrate following binge alcohol drinking, resulting from ethanol-induced PKC-activation that hyperphosphorylates GSK 3β to cause enhanced calcineurin-NFAT-Csx/Nkx2.5 signaling. These observations elucidate for the first time the potential mechanisms underlying the clinically well-recognized, but mechanistically enigmatic, “holiday heart syndrome”.

Key Words: Alcohol; Atrial fibrillation; Electrophysiology; Ion channels

Atrial fibrillation (AF) is the most commonly sustained cardiac arrhythmia. Although certain risk factors, such as age, hypertension, serum metabolites and heart failure are well-established,^{1–3} the causes and mechanisms remain unknown in many patients. There is conflicting evidence regarding an association between long-term alcohol consumption and the risk of AF.⁴ In contrast, an association between binge alcohol-drinking and AF, often termed “holiday heart syndrome”, has been widely recognized.⁵ Holiday heart syndrome refers to cardiac arrhythmias, particularly AF, that occur after an alcoholic binge in individuals showing no other evidence of heart disease, which usually convert to normal sinus rhythm

Editorial p????

within 24 h.⁶ Several potential mechanisms have been postulated by which an alcoholic binge could cause arrhythmias. Alcohol-induced increases in plasma-free fatty acids and catecholamines are thought to be arrhythmogenic,⁷ as is the principal metabolite of alcohol, acetaldehyde.⁸ However, the precise electrophysiological mechanisms that connect acute ethanol consumption to AF are largely unknown.

Pulmonary veins (PVs) are important foci for initiation of paroxysmal AF and are also associated with AF maintenance.⁹ PVs contain cardiomyocytes with electrical activ-

Received March 31, 2020; revised manuscript received July 20, 2020; accepted August 7, 2020; J-STAGE Advance Publication released online October 7, 2020 Time for primary review: 20 days

Department of Pathophysiology (Y.W., M.M., D.L., K.O.), Department of Clinical Examination and Diagnostics (Y.W., N.T., T.S., K.O.), Oita University School of Medicine, Yufu, Japan; Montreal Heart Institute Research Center, University of Montreal, Montreal (S.N.), Canada

The first two authors contributed equally to this study (Y.W., M.M.).

Mailing address: Katsushige Ono, MD, PhD, Department of Pathophysiology, Oita University School of Medicine, 1-1 Idaigaoka, Hasama, Yufu 879-5593, Japan. E-mail: ono@oita-u.ac.jp

All rights are reserved to the Japanese Circulation Society. For permissions, please e-mail: cj@j-circ.or.jp

ISSN-1346-9843

